Review Article

Biomarkers in dentin pulp complex

Shaik Ali Hassan^{1,*}, Sumit Bhateja¹, Geetika Arora², Francis Prathyusha³¹Dept. of Dental, Manav Rachna Dental College, Faridabad, Haryana, India²Inderprastha Dental College & Hospital, Ghaziabad, Uttar Pradesh, India³Dr. Francis Maxillofacial and Dental Clinic, India

ARTICLE INFO

Article history:

Received 05-06-2020

Accepted 16-06-2020

Available online 18-08-2020

Keywords:

Biomarkers

Periapical

BSP

MMP

Pulp.

ABSTRACT

Biomarkers are functional elements at the level of cells or molecules, play an important role in the health and disease. Tooth dentin-pulp complex houses several biomarkers at different stages of development, and the lack of these biomarkers yield developmental disorders. Furthermore, biomarkers play a very important role in the pathogenesis of dental caries, pulp and periapical pathoses in two ways - they is an important element in the process of detection of pathological and they assist in the accurate diagnosis of pathological conditions.

© 2020 Published by Innovative Publication. This is an open access article under the CC BY-NC license (<https://creativecommons.org/licenses/by-nc/4.0/>)

1. Introduction

The term "biomarker" is created from a portmanteau of "Biological markers" which refers to broad subcategory of medical signs that can be measured accurately and reproducibly.^{1,2} National Institutes of Health Biomarkers the Working Group has defined the definition of a biomarker as a characteristic that is objectively measured and evaluated as an indicator of normal biological processes, pathogenic processes, or pharmacologic responses to a therapeutic intervention.³ Basically, a biomarker can be considered as a response functional, psychological, biochemical or cellular level molecular interactions. Biomarkers are being widely used in research and has proven to be a boon in the medical field to assist in diagnosis and treatment.

2. Various Biomarkers

The dentin and pulp together form the dentin-pulp complex. The outermost cell layer at the boundary of pulp tissue and predentin consists of the odontoblasts, the cells that form

and mineralize the dentin matrix⁴

3. Osteocalcin

Odontoblasts express a glycoprotein called osteocalcin (OCN) in the matrix of the dentine.⁵ He is known to be one of reparative molecules and expression occurs in response to injury of the dental pulp. It is believed to be collated with collagen fibers and is also found embedded in the tertiary dentin, where its expression occurs in response to the cavity preparation.⁶ In addition, it is also considered as a marker for mature osteoblasts and has been used as markers of osteoblast-like differentiation mineralization odontoblast / on Stem cells from dental pulp.⁷

OCN is expressed before the start of the mineralization and is the highest non-collagenous protein in bone extracellular matrix. It is suggested to be essential in interceding.^{7,8} osteoclast differentiation. OCN was found associated with proteins such as macrophage granulocyte macrophage colony-stimulating factor (GM-CSF) in cells forming bone. Expression NCBs occurs in the terminal differentiation of macrophages osteoblasts. It is expressed

* Corresponding author.

E-mail address: alishaikhassan@gmail.com (S. A. Hassan).

in response to the preparation of the cavity associated with the collagen fibers and is also found in the tertiary dentin.⁹

4. Osteonectin

It happens to be a major non-collagen matrix protein bone and dentin.¹⁰ odontoblasts and beef, and odontoblasts predentine in human prenatal and postnatal samples were exposed. Its various roles in the initiation of mineralization was.¹¹ shown Because of their study, ON is indicated as a protein associated with the formation of collagen in mineralized tissues, such as bone and human dentin.

5. Dentin sialophosphoprotein

DSPP is the initial translation product DSPP envoy RNA (mRNA) which is then cleaved to the teeth and phosphoprotein dentin sialoprotein (DSP).^{12–14} That primitive is considered to be Special dentin until detected on bone.¹⁵ However, the expression of dentin is about 400 times that of bone. Being one of the non-collagenous proteins necessary foundation in the development of teeth and mineralization, it is mainly expressed in odontoblasts. DSPP role in dentinogenesis has been well demonstrated.¹⁵ (Qin et al., 2002), so it remains a major marker for odontoblast differentiation. DSPP, osteocalcin and matrix extracellular phosphoglycoprotein (MEPE) grew up in a time expression Depending on the dental pulp cells (DPC) induced.^{13,16–18} culture.

6. Thyrotropin-Releasing Hormone -Degrading Enzyme

This is known as pyroglutamyl peptidase II and shows the absolute functional specificity for substrates, TRH.^{19,20} It is oriented extracellular, a membrane-associated peptidase (ectopeptidase) and a function to stop the peptide-mediated cell signaling. TRH-DE has been found in the dental pulp by microarray analysis and real time RT-PCR analysis of.²⁰ TRH-DE mRNA expression in dental pulp stem cells / progenitor (CD105 + and CD31- Side population (SP) cells were enhanced by induced nerve cell. Its presence in the process of neuronal The dental pulp is confirmed by immunohistochemistry and in situ hybridization. In addition, the TRH-DE mRNA was expressed in pulp regeneration 28 days after transplant CD31- SP cells into a root canal after pulpectomy.

7. Matrix metalloproteinase

MMPs are synthesized by cells of connective tissue such as fibroblasts, osteoblasts, and odontoblasts and secreted into the extracellular matrix. Several matrix metalloproteinases (MMP) have been identified in the dentin and pulp by polymerase chain reaction (PCR) and immunohistochemistry.^{21–23} MMPs have been

classified into six groups based on their structural homology and their substrate specificity as collagenase (MMP-1, MMP-8, MMP-13 and MMP-18), gelatinases (MMP-2 and MMP-9), stromelysins (MMP-3, MMP-10 and MMP-11), MMP transmembrane or MT-MMP (MMP-14, MMP-15, MMP-16, MMP-17, MMP-24 and MMP-25), and others (MMP-12, MMP-19, MMP-20, MMP-21, MMP-22, MMP-23, MMP-27 and MMP-28).²²

Some MMPs have been identified in human mineralization dentin, ie collagenase MMP-8²⁴ gelatinases (25) and enamelysin.²³ MMP MMP-20 is present in the saliva and the dentin is host-derived and activated at acidic pH resulting from the release of lactate of cariogenic bacteria.^{25,26} At neutral pH, the effect of MMP in remodeling of the extracellular matrix. tissue damage always colligated with aberrant expression of MMP. MMP role in various physiological processes in the dentin-pulp complex has good roles including organizational understood. The matrix prior to mineralization, the mineralization controls, peritubular dentin formation, and changes in the matrix during aging.²⁷

8. Bone sialoprotein

BSP has been contained within the mineralized dentin and modulation of MMP-2. use the collected RNA was isolated from the pulp tissue of both health and caries, it has been observed that there is 8-fold up-regulation in gene expression in the pulp BSP teeth with active caries.²⁸ However, the expression of the pulp tissue MMP-2 in response to the caries process remains unknown. Immunohistochemical analysis of extracted third molars and premolars did not reveal any detection of MMP-2 and BSP in tubular lumen of sound dentin. Therefore, both BSP and MMP-2 may involved in host defense mechanisms that lead to calcification areas affected by caries.

9. Biomarkers in dental pulp complex in diseases

Infection, exposure, trauma and chemicals may result in losses pulp vitality. As a result of these tooth injury, the signal for progenitor stem cells were sent to stimulate the differentiation, proliferation, and migration as part of reparative dentinogenesis. Influx and macrophage recruitment of polymorphonuclear (PMN) is the beginning In response to bacteria and their metabolites. In addition to fibroblast proliferation and angiogenesis, macrophage infiltration, lymphocytes and plasma cells are a typical feature in more chronic conditions. MMPs have also been identified for both pulp and periapical.^{29–31} inflammation. This group of enzymes responsible for degradation of extracellular matrix (ECM), as seen in inflammationpulp and periapical region. The destruction of the ECM generated by component of intracellular bacteria, bacterial metabolites, and other molecules, leading to the formation

of periradicular lesions.³²

Basically, MMP is a group of structurally related but endopeptidases expressed in genetically distinct normal tissues at low levels, but regulated inflammation. That also interesting to note that the levels of MMP-8, which is usually higher in the case of periapical exudate, reduced significantly after the first visit root canal treatment.³³ IHC staining showed MMP-8 in the pulp and periapical granulomas with PMN become larger cell types to express MMP-8.

Enzyme linked immunosorbent assay has shown that MMP-1 levels are below the limit of detection in healthy and inflamed pulp, while the level of MMP-9 was significantly increased in inflamed human dental pulp. In contrast, levels of MMP-2 and MMP-3 reduced the inflamed pulp symptoms than with a normal pulp.³⁴ This suggests a major role of MMP-9 in the degradation of inflamed human dental pulp tissue. Role MMP-9 in the inflamed pulp was then supported by another report.³⁵ Another study also noted that MMP-9 mRNA gene is increased in the inflamed pulp.³⁴ Furthermore, in-situ localization of MMP-9 expression in specimens showed significantly pulp higher expression of MMP-9 in the inflamed pulp compared to clinically healthy pulp. Increased activity of MMP-9 and MMP-2 has also has been shown in the gingival sulcus fluid teeth with periapical lesions.³⁵ Thus, the use of MMP-9 and MMP-2 as a biological markers can be proved. Furthermore, periapical granuloma showed higher MMP-9 and MMP-13 activity compared with³⁴ radicular cysts. Osteocalcin role in the pulp pathoses not clearly explained until recently, when it was shown that the expression of osteocalcin was higher in comparison with irreversible reversible pulpitis pulpitis,⁸ Osteocalcin is reparative molecules in the dental pulp and in terms of improvement, it is localized in the cell and the surrounding matrix areas of calcification in the cells surrounding the blood vessels. However it's not in normal tissue. These findings correlated positively with angiogenic markers such as vascular endothelial growth factor (VEGF) and fibroblast growth factor (FGF), supporting role in the repair of pulp.

10. Biomarkers as diagnostic tool

As there is a significant expression of MMPs pulp and pathological inflammation, MMP analysis of periapical exudate can an option to demonstrate and monitor the activity of inflammatory and as an indicator of the success of root canal treatment that has previous periapical lesions. There is a strong denotation that MMP-8 periapical inflammation appear on the site because it is still on display in second visit root canal treatment regardless of the deletion pulp networks and canals cleaned during the first appointment that implies that the active phase of periapical inflammation sites still exist. The loss of partial onset inflammation and healing is reflected by the lack of MMP-8

in exudates during root canal third visit. Therefore, the level of MMP-8 could serve as Biochemical indicators to assess the inflammatory status of the periapical tissues. MMP measurement of the root canal during treatment serves as a potential prospect as a diagnostic tool to assess periapical inflammatory conditions. MMP-9 also serves as potentially unreliable marker as levels significantly improve in pulp inflammation.³⁵ The presence of molecular differential expression in the cyst or granuloma has the potential to provide information to distinguish between periapical cysts of granuloma before executing Endodontic procedures as cysts have a lower healing rate.³⁶ Zehnder et al is the first to provide clinical relevance for the use of MMP-9 as a marker for the pulp pathoses.^{37–39} They performed a study to improve the diagnosis of the pulp to assess the level of MMP-9 in dentin fluid diagnosed with irreversible symptoms pulpitis and healthy peers. In order to perform non-invasive assessment of pulp, dentin fluid collection of dentin wounds can be done after the cavity preparation. After access cavity is prepared, sterile folded polyvinylidene difluoride (PVDF) membrane filters used to collect fluid dentin of dentin open and the fluid is then charged MMP-9 assay neon. This particular assay has been claimed has a sensitivity that is superior to conventional ELISA for functions based substrate turnover effect in the liberation of the fluorophore. In conclusion, dentin fluid samples from dental symptoms had significantly higher MMP-9 levels compared for healthy pulp.

11. Source of Funding

None.

12. Conflict of Interest

None.

References

1. Scaros O, Fislser R. Biomarker technology roundup: from discovery to clinical applications, a broad set of tools is required to translate from the lab to the clinic. *BioTech*. 2005;38(4S):S30–2.
2. Strimbu K, Tavel JA. What are biomarkers? *Curr Opin HIV AIDS* . 2010;5(6):463–6.
3. Biomarkers and surrogate endpoints: preferred definitions and conceptual framework. *Clin Pharmacol Therapeutics*. 2001;69:89–95.
4. Cate T, R A. Structure of the Oral Tissues. Mosby; 1994. p. 45–57.
5. Abd-Elmeguid A, Abdeldayem M, Kline LW, Moqbel R, Vliagoftis H, Yu DC, et al. Osteocalcin Expression in Pulp Inflammation. *J Endod*. 2013;39(7):865–72.
6. Hirata M, Yamaza T, Mei YF, Akamine A. Expression of osteocalcin and Jun D in the early period during reactionary dentin formation after tooth preparation in rat molars. *Cell Tissue Res*. 2005;319(3):455–65.
7. de Vries IG, Quartier E, Boute P, Wisse E, Coomans D. Immunocytochemical Localization of Osteocalcin in Developing Rat Teeth. *J Dent Res*. 1987;66(3):784–90.
8. Ishida M, Amano S. Osteocalcin fragment in bone matrix enhances osteoclast maturation at a late stage of osteoclast differentiation. *J Bone Mineral Metab*. 2004;22(5):415–29.

9. Evans DB, Bunning RAD, Russell RGG. The effects of recombinant human granulocyte-macrophage colony-stimulating factor (rhGM-CSF) on human osteoblast-like cells. *Biochem Biophys Res Commun*. 1989;160(2):588–95.
10. Fujisawa R, Kuboki Y. Changes in levels of osteonectin in bovine dentine during tooth development. *Arch Oral Biol*. 1989;34(2):89–92.
11. Reichert T, Störkel S, Becker K, Fisher LW. The role of osteonectin in human tooth development: An immunohistological study. *Calcif Tissue Int*. 1992;50(5):468–72.
12. Butler WT. Dentin specific proteins. *Methods Enzymol*. 1987;145:290–303.
13. Butler WT, Ritchie H. The nature and functional significance of dentin extracellular matrix proteins. *Int J Dev Biol*. 1995;39:169–79.
14. Butler WT, Brunn JC, Qin C, Mckee MD. Extracellular matrix proteins and the dynamics of dentin formation. *Connect Tissue Res*. 2002;43:301–7.
15. Qin C, Brunn JC, Cadena E, Ridall A, Tsujigiwa H, Nagatsuka H, et al. The Expression of Dentin Sialophosphoprotein Gene in Bone. *J Dent Res*. 2002;81(6):392–4.
16. Sreenath T, Thyagarajan T, Hall B. Dentin sialophosphoprotein knockout mouse teeth display widened predentin zone and develop defective dentin mineralization similar to human dentinogenesis imperfect type III. *J Biol Chem*. 2003;278:24874–80.
17. Zhang X, Zhao J, Li C. DSPP mutation in dentinogenesis imperfect Shields type II. *Nat Genet*. 2001;27:151–2.
18. Papagerakis P, Berdal A, Mesbah M, Peuchmaur M, Malaval L, Nydegger J, et al. Investigation of osteocalcin, osteonectin, and dentin sialophosphoprotein in developing human teeth. *Bone*. 2002;30(2):377–85.
19. BAUER K, CARMELIET P, SCHULZ M, BAES M, DENEFC. Regulation and Cellular Localization of the Membrane Bound Thyrotropin-Releasing Hormone-Degrading Enzyme in Primary Cultures of Neuronal, Glial and Adenohypophyseal Cells*. *Endocrinol*. 1990;127(3):1224–33.
20. Yamamoto T, Murakami M, Ishizaka R, Iohara K. Identification of Thyrotropin-Releasing Hormone (TRH)-Degrading Enzyme as a Biomarker for Dental Pulp Tissue. *Dent*. 2012;02(01):114.
21. Palosaari H, Pennington CJ, Larmas M, Edwards DR, Tjäderhane L, Salo T. Expression profile of matrix metalloproteinases (MMPs) and tissue inhibitors of MMPs in mature human odontoblasts and pulp tissue. *Eur J Oral Sci*. 2003;111(2):117–27.
22. Chaussain-Miller C, Fioretti F, Goldberg M, Menashi S. The Role of Matrix Metalloproteinases (MMPs) in Human Caries. *J Dent Res*. 2006;85(1):22–32.
23. Sulkala M, Larmas M, Sorsa T, Salo T, Tjäderhane L. The Localization of Matrix Metalloproteinase-20 (MMP-20, Enamelysin) in Mature Human Teeth. *J Dent Res*. 2002;81(9):603–7.
24. Sulkala M, Tervahartiala T, Sorsa T, Larmas M, Salo T, Tjäderhane L, et al. Matrix metalloproteinase-8 (MMP-8) is the major collagenase in human dentin. *Arch Oral Biol*. 2007;52(2):121–7.
25. Mazzoni A, Pashley D, Nishitani Y, et al. Reactivation of inactivated endogenous proteolytic activities in phosphoric acid-etched dentine by etch-and-rinse adhesives. *Biomater*. 2006;27(25):4470–6.
26. Mazzoni A, Pashley DH, Tay FR. Immunohistochemical identification of MMP-2 and MMP-9 in human dentine: correlative FEI-SEM/TEM analysis. *J Biomed Mater Res*. 2009;88:697–703.
27. Boushell LW, Kaku M, Mochida Y, Bagnell R, Yamauchi M. Immunohistochemical localization of matrixmetalloproteinase-2 in human coronal dentin. *Arch Oral Biol*. 2008;53(2):109–16.
28. McLachlan JL, Smith AJ, Bujalska IJ, Cooper PR. Gene expression profiling of pulpal tissue reveals the molecular complexity of dental caries. *Biochim Biophys Acta*. 2005;1741(3):271–81.
29. Gusman H, Santana RB, Zehnder M. Matrix metalloproteinase levels and gelatinolytic activity in clinically healthy and inflamed human dental pulps. *Eur J Oral Sci*. 2002;110(5):353–7.
30. Tsai CH, Chen YJ, Huang FM, Su YF, Chang YC. The upregulation of matrix metalloproteinase-9 in inflamed human dental pulps. *J Endod*. 2005;31:860–2.
31. Paula-Silva FWD, Silva NJD, Silva LAD. High matrix metalloproteinase activity is a hallmark of periapical granulomas. *J Endod*. 2009;35:1234–42.
32. Yang W, Harris MA, Cui Y, Mishina Y, Harris SE, Gluhak-Heinrich J, et al. Bmp2 Is Required for Odontoblast Differentiation and Pulp Vasculogenesis. *J Dent Res*. 2012;91(1):58–64.
33. Boushell LW, Nagaoka H, Nagaoka H, Yamauchi M. Increased Matrix Metalloproteinase-2 and Bone Sialoprotein Response to Human Coronal Caries. *Caries Res*. 2011;45(5):453–9.
34. Paula-Silva FWD, Silva D, Silva NJD, A L. High matrix metalloproteinase activity is a hallmark of periapical granulomas. *J Endod*. 2009;35:1234–1242.
35. Belmar MJ, Pabst C, Martínez B, Hernández M. Gelatinolytic activity in gingival crevicular fluid from teeth with periapical lesions. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2008;105:801–6.
36. Karapanou V, Kempuraj D, Theoharides TC. Interleukin-8 Is Increased in Gingival Crevicular Fluid from Patients with Acute Pulpitis. *J Endod*. 2008;34(2):148–51.
37. Burgener B, Ford AR, Situ H, Fayad MI, Hao JJ, Wenckus CS, et al. Biologic Markers for Odontogenic Periradicular Periodontitis. *J Endod*. 2010;36(8):1307–10.
38. Zehnder M, Wegehaupt FJ, Attin T. A First Study on the Usefulness of Matrix Metalloproteinase 9 from Dentinal Fluid to Indicate Pulp Inflammation. *J Endod*. 2011;37(1):17–20.
39. Rauschenberger CR, McClanahan SB, Pederson ED, Turner DW, Kaminski EJ. Comparison of human polymorphonuclear neutrophil elastase, polymorphonuclear neutrophil cathepsin-G, and α 2-macroglobulin levels in healthy and inflamed dental pulps. *J Endod*. 1994;20(11):546–50.

Author biography

Shaik Ali Hassan Dental Surgeon

Sumit Bhateja HOD

Geetika Arora Reader

Francis Prathyusha BDS

Cite this article: Hassan SA, Bhateja S, Arora G, Prathyusha F. **Biomarkers in dentin pulp complex.** *IP J Paediatr Nurs Sci* 2020;3(2):34-37.